


# Coherent Matter Travelling Wave Beams

And their possible role in making 'Strange Radiation' tracks - Part 2










#### Plan


- Discuss methods of coherent matter production
- Behaviour of 'balls of fire' on and in metals
- Behaviour of observed traces vs witness marks
- Product from a dead 'Coherent'
   Matter Travelling Wave Beam'
- Special Treat!

#### OHMA, 2019












Matsumoto, 1993, Journal of Fusion Technology, 23 (1), 193-113





# Links to learn about Strange Radiation



Lutz Jaitner, focussed on geometry and history

http://condensed-plasmoids.com/history.html

http://condensed-plasmoids.com/condensed plasmoids lenr.pdf

Edward Lewis, first to tell Matsumoto, good focus on health impacts

http://www.scientificrevolutions.com/

http://tc38.metawerx.com.au/oldsite/2021/lewishealth2021summer.pdf

Bob Greenyer - Amazing tracks (Alexander Parkhomov)

https://youtu.be/cUul-6yp8G8



# Slobodan Stankovic

Shown ICCF-22 September 2019


# Anatoly Klimov - Plasmatron 17 June 2020

# VEGA Various Experiments in Glowing Action

- Team is Henk (Holland), Dave (Canada) and myself
- Use of ganged microwave transformers,
   to produce several 100V DC for discharge
- Simple vacuum chambers, not high vacuum
- Residual air with gasses such as Ar and H

### VEGA Nearly stable

HENK/MFMP











- Ignore cathode/anode
- Ignore pressure waves
- Interact with dense material
- Large changes in direction and speed
- Split/merge with divided segments showing same in-phase structure
- Interact at a distance and in anticipation of interaction







# Some nice tracks





# Going through metal?







## Natural Ball lighting traces vs CMTW





Ref






Coherent
matter travelling
waves in VEGA

Trace 1A scaled 3x and rotated

Trace 1B scaled 3x and rotated




Constrained 'Strange Radiation' result presented at the EADS colloqium in 2010 by Claude Daviau, Didier Priem and Guillaume Racineux, observed in Nante when replicating the exploding Titanium approach of Leonid Urutskoev. Published 2015, J-F Geneste




3 dimensional trace 1A scaled, rotated and distorted to 'modify frequency'











# Thankyou - Q&A

- To the organisers of ICMNS 2021
- Alan Goldwater and MagicSoundLabs
- To all of the crowd researchers that made this possible
- To the generous donors, in particular to Charles and Sho that made this trip possible
- Henk, Dave, Slobodan, John, Peter and all the many experimentalists working together openly to solve this problem